



US006080553A

United States Patent [19]  
Sogabe et al.

[11] Patent Number: 6,080,553  
[45] Date of Patent: \*Jun. 27, 2000

[54] CREATINE AMIDINOHYDROLASE,  
PRODUCTION THEREOF AND USE  
THEREOF

3,907,644 9/1975 Mollering et al. .... 195/99  
5,451,520 9/1995 Furukawa et al. .... 435/227

[75] Inventors: Atsushi Sogabe; Takashi Hattori;  
Yoshiaki Nishiyama; Yoshihisa  
Kawamura, all of Tsuruga, Japan

## FOREIGN PATENT DOCUMENTS

62 091182 4/1987 Japan  
07 265074 10/1995 Japan

[73] Assignee: Toyo Boseki Kabushiki Kaisha,  
Osaka, Japan

[\*] Notice: This patent issued on a continued prosecution application filed under 37 CFR 1.53(d), and is subject to the twenty year patent term provisions of 35 U.S.C. 154(a)(2).

[21] Appl. No.: 08/799,897

[22] Filed: Feb. 13, 1997

## [30] Foreign Application Priority Data

Feb. 13, 1996 [JP] Japan 8-025435

[51] Int. Cl.<sup>7</sup> C12Q 1/34; C12N 9/78;  
C12N 1/20; C12N 1/00

[52] U.S. Cl. 435/18; 435/227; 435/192;  
435/252.3; 435/320.1; 435/829; 435/252.33

[58] Field of Search 435/18, 227, 252.1,  
435/320.1, 829, 192, 252.3, 252.33

## [56] References Cited

## U.S. PATENT DOCUMENTS

3,806,420 4/1974 Holz et al. 195/66

Primary Examiner—Elizabeth Slobodyansky  
Attorney, Agent, or Firm—Leydig, Voit & Mayer, Ltd.

## [57] ABSTRACT

A creatine amidinohydrolase having the following physico-chemical properties:

Action: catalyzing the following reaction;  
creatine+H<sub>2</sub>O→sarcosine+urea

Optimum temperature: about 40–50° C.

Optimum pH: pH about 8.0–9.0

Heat stability: not more than about 50° C. (pH 7.5, 30 min)

Km value for creatine in a coupling assay using a sarcosine oxidase and a peroxidase: about 3.5–10.0 mM

Molecular weight: about 43,000 (SDS-PAGE)

Isoelectric point: [3.5] 4.5

a method for producing said enzyme, comprising culture of microorganism producing said enzyme, a method for the determination of creatine or creatinine in a sample using said enzyme, and a reagent therefor.

23 Claims, 2 Drawing Sheets